

Pop III Hypernova Nucleosynthesis and Abundances in Very Metal-Poor Halo Stars

Hideyuki Umeda¹, Ken'ichi Nomoto¹

¹ *Department of Astronomy, School of Science, The University of Tokyo,
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*

Abstract

We calculate evolution and nucleosynthesis in massive Pop III stars with $M = 13 \sim 270M_{\odot}$, and compare the results with abundances of very metal-poor halo stars. The observed abundances can be explained by the energetic core-collapse supernovae with $M \lesssim 130M_{\odot}$ (“hypernovae”) but not by pair-instability supernovae (PISNe) with $M \sim 140 - 270M_{\odot}$. This result constrain the IMF for the Pop III and very metal-poor Pop II stars.

Observed Abundance and Hypernova Nucleosynthesis

The observed abundances of metal-poor halo stars show quite interesting trends. There are significant differences between the abundance patterns in the iron-peak elements below and above $[\text{Fe}/\text{H}] \sim -2.5$. For $[\text{Fe}/\text{H}] \lesssim -2.5$, the mean values of $[\text{Cr}/\text{Fe}]$ and $[\text{Mn}/\text{Fe}]$ decrease toward smaller metallicity, while $[\text{Co}/\text{Fe}]$ and $[\text{Zn}/\text{Fe}]$ increases (McWilliam et al. 1995; Ryan et al. 1996; Primas et al. 2000; Blake et al. 2001; see also Fig.4). The Galaxy was not well mixed in such a early stage, and the abundance pattern of each supernova (SN) may be kept in the very metal-poor stars. However, these trend could not be explained with the previous results of nucleosynthesis by core-collapse SNe such as Woosley & Weaver (1995), Nomoto et al. (1997) and Limongi et al. (2000). Therefore, we need reconsideration for the nucleosynthesis of very metal-poor stars.

We found that these trends are explained simultaneously in the context of Fe core-collapse SNe, if the mass from complete Si-burning is relatively large compared with that from incomplete Si-burning (Nakamura et al. 1999). However, all previous model calculations including Nakamura et al. (1999) underproduced the Zn/Fe ratio significantly (Fig. 1 left). In Umeda & Nomoto (2002, UN02 hereafter), we find that $[\text{Zn}/\text{Fe}]$ is larger for deeper mass-cuts, smaller neutron excess, and larger explosion energies. Among them the large explosion energy is most important to realize the large $[\text{Zn}/\text{Fe}]$ ratio (Fig. 1 right). The observed trends of the abundance ratios among the iron-peak elements are better explained with this high energy (“Hypernova”) models than the simple “deep” mass-cut effect, because the overabundance of Ni can be avoided in the hypernova models.

Pair Instability Supernovae

We also investigate the yields of pair-instability supernova explosions of $M \simeq 140 - 300M_{\odot}$ stars. In Fig.2 we compare the abundance pattern of a hypernova and a PISN model. As can be seen $[\text{Zn}/\text{Fe}]$ of PISNe are always small, because relatively large incomplete Si-burning region is formed in the explosion (UN02). Therefore, the abundance features of very metal-poor stars cannot be explained by pair-instability supernovae.




Figure 1: Observed range of $[Zn/Fe]$ for the very metal-poor stars ($[Fe/H] \lesssim -3$) compared with the yields of theoretical models with low explosion energy $E_{51} = E_{\text{exp}}/10^{51} \text{ erg} = 1$ (left) and with high energies $E_{51} = 2 \sim 50$ (right panel). The observed large $[Zn/Fe]$ value and also other unexpected trends in $[Co, Mn, Cr/Fe]$ can be explained with this high energy (“Hypernova”) models.

$[Fe/H]$ and SNe Mass, Energy

We have discussed that the large $[Zn/Fe]$ ratio can be explained with hypernova nucleosynthesis of $M \lesssim 130M_{\odot}$ stars. However, in order to explain the observed trend for $[Fe/H]$, it is necessary to explain why hypernova contribution is dominant for lower $[Fe/H]$. Suppose that the star-formation is induced by the supernova shock. Then the $[Fe/H]$ of the next generation stars is determined by the ratio of Fe mass synthesized by the SN and the hydrogen mass swept by the supernova shock. Since the latter is inversely proportional to the explosion energy E , the following relation is obtained (Ryan et al. 1996; Shigeyama & Tsujimoto 1998):

$$Fe/H \propto M(^{56}\text{Ni})/E. \quad (1)$$

In Fig.3 we show the similar relation for Mg as a function of the mass and energy of SNe. Although the ejected Fe mass is theoretically uncertain, in most of the observed stars, Mg/Fe ratio is roughly the same and typically $[Mg/Fe] \simeq 0.4$. Thus $[Mg/H] \sim [Fe/H] + 0.4$. As shown in this figure, if the explosion energy E_{51} is fixed, $[(Mg, Fe)/H]$ increases with mass, because more Mg is ejected for more massive stars. However the energy factor is much more important, and hence massive hypernova models have smaller $[(Mg, Fe)/H]$ than less massive normal SNe II. In this way, hypernova like nucleosynthesis patterns can be seen in the very metal-poor stars.

Finally in Fig.4, together with the observation points, we plot the abundance pattern of Fe-peak elements for a normal SNe II ($20M_{\odot}$, $E_{51} = 1$) and a hypernova ($25M_{\odot}$, $E_{51} = 30$) models in UN02. Here the relative position in $[Fe/H]$ is determined by the relation (1). Although there are some inconsistencies between the model and observations, we can explain the trend in the observation. As discussed in UN02, the Mn abundance is very sensitive to Y_e of the progenitor, and thus the enhancement of factor 7 is not so difficult to realize. The remaining problem is the overproduction of Cr and underproduction of Co, which should be investigated further in future work.




Figure 2: The abundance pattern of a hypernova (left) and a PISN model (right panel), normalize to the solar abundance. For PISNe with initial masses $M \simeq 140 - 300M_{\odot}$ $[\text{Zn}/\text{Fe}]$ is small, so that the abundance features of very metal-poor halo stars cannot be explained by these SNe.




Figure 3: $[(\text{Mg}, \text{Fe})/\text{H}]$ determined by the relation (1) as a function of SN mass and energy. The constant C is chosen to normalize the $(20M_{\odot}, E_{51} = 1)$ case to be $[\text{Mg}/\text{H}] = -2$.

References

- [1] Blake, L.A.J., Ryan S.G., Norris, J.E., & Beers, T.C. 2001, Nucl.Phys.A., 688, 502
- [2] Limongi, M., Straniero, O., & Chieffi, A. 2000, ApJS, 129, 625
- [3] McWilliam, A., Preston, G. W., Sneden, C., & Searle, L. 1995, AJ, 109, 2757
- [4] Nakamura, T., Umeda, H., Nomoto, K., Thielemann, F.-K., & Burrows, A. 1999, ApJ, 517, 193
- [5] Nomoto, K., Hashimoto, M., Tsujimoto, T., Thielemann, F.-K., Kishimoto, N., Kubo, Y., & Nakasato, N. 1997, Nucl. Phys. A616, 79
- [6] Primas, F., Reimers, D., Wisotzki, L., Reetz, J., Gehren, T., & Beers, T.C. 2000, in The First Stars, ed. A. Weiss, T. Abel, & V. Hill (Berlin: Springer), 51




Figure 4: Observational points of Fe-peak elements (small symbols) and the theoretical abundance patterns for a normal SNe II ($20M_{\odot}$, $E_{51} = 1$) and a hypernova ($25M_{\odot}$, $E_{51} = 30$) models in UN2002 (large open circles). Here, Mn abundance is multiplied by 7.

- [7] Ryan, S. G., Norris, J. E., & Beers, T. C. 1996, ApJ, 471, 254
- [8] Shigeyama, T., & Tsujimoto, T. 1998, ApJ, 507, L135
- [9] Umeda, H., Nomoto, K. (2002), ApJ, 565, 385 (UN02)
- [10] Woosley, S.E., & Weaver, T.A. 1995, ApJS, 101, 181